


# A typical journey from the onset of symptoms to diagnosis



3 in 4 were told their symptoms weren't cardiac

75%



Over half (59%) of respondents called an ambulance due to symptoms of INOCA



2 in 3 were discharged without treatment from A&E

67%



Only 8 out 100 ambulance crews understood INOCA

## Misdiagnoses

1 in 4 respondents were prescribed antidepressants

42%



1 in 3 respondents were referred to a Psychiatrist

32%



1 in 2 respondents were told that symptoms were GERD

54%



6% were diagnosed on first consultation

29% had 2 – 3 consultations

Consultations prior to diagnosis

35% had 4 – 6 consultations

Number of cardiologists consulted

20%



26%



50%



25% had over 6 consultations

49%

Just under half of respondents are currently under the care of an INOCA specialist

Martha Gulati, Najah Khan, Maria George, Colin Berry, Alaide Chieffo, Paolo G. Camici, Filippo Crea, Juan-Carlos Kaski, Mario Marzilli, C. Noel Bairey Merz

[doi.org/10.1016/j.ijcard.2022.09.047](https://doi.org/10.1016/j.ijcard.2022.09.047)

Posters by Alex Koch

[www.INOCAInternational.com](http://www.INOCAInternational.com)

Scan for paper

